

**SEMMELWEIS EGYETEM  
DOKTORI ISKOLA**

**Ph.D. értekezések**

**2954.**

**EKE CSABA**

**Szív-és érrendszeri betegségek élettana és klinikuma  
című program**

Programvezető: Dr. Merkely Béla, egyetemi tanár

Témavezető: Dr. Székely Andrea, egyetemi tanár

# **Association between hepatic venous flow and adverse outcomes after cardiac surgery**

**PhD thesis**

**Csaba Eke MD**

Doctoral School of Theoretical and Translational Medicine

Semmelweis University



Supervisor:

Székely Andrea, MD, DSc

Official reviewers:

Fülöp András MD, PhD

Babik Barna MD, PhD

Head of the Complex Examination Committee:

Karádi István MD, DSc

Members of the Complex Examination Committee:

Farkas Henriette MD, DSc

Charaf Hassan DSc.

Budapest  
2023



Table of contents

<b>1. Table of contents</b> .....	<b>4</b>
<b>3. List of abbreviations</b> .....	<b>6</b>
<b>4. Introduction</b> .....	<b>8</b>
1. <i>Fluid balance and fluid overload in critically ill patients</i> .....	8
2. <i>Hepatic venous flow and abdominal congestion</i> .....	10
3. <i>Postoperative right ventricle dysfunction, congestion, and acute kidney injury after cardiac surgery</i> .....	15
<b>5. Objectives</b> .....	<b>19</b>
<b>6. Methods</b> .....	<b>20</b>
1. <i>Structure of the studies</i> .....	20
2. <i>Ultrasound examinations</i> .....	22
3. <i>Main goals</i> .....	24
4. <i>Statistical analysis</i> .....	24
<b>7. Results</b> .....	<b>26</b>
1. <i>Connection between preoperative hepatic venous profile and postoperative acute kidney injury (study A)</i> .....	26
1. <i>Descriptive statistics</i> .....	26
2. <i>Univariable and multivariable regression</i> .....	29
2. <i>Connection between hepatic venous congestion and morbidity (study B)</i> .....	33
1. <i>Descriptive statistics</i> .....	33
2. <i>Regressions</i> .....	38
<b>8. Discussion</b> .....	<b>40</b>
1. <i>Perioperative fluid overload</i> .....	40

2.	<i>Mechanical ventilation</i> .....	41
3.	<i>Echocardiography</i> .....	42
4.	<i>Acute kidney injury</i> .....	42
5.	<i>Other liver parameters</i> .....	43
6.	<i>Limitations</i> .....	44
<b>9.</b>	<b>Conclusions</b> .....	<b>45</b>
<b>10.</b>	<b>Summary</b> .....	<b>46</b>
<b>11.</b>	<b>References</b> .....	<b>48</b>
<b>12.</b>	<b>Bibliography of the candidate's publications</b> .....	<b>60</b>
<b>13.</b>	<b>Acknowledgements</b> .....	<b>63</b>

## 2. List of abbreviations

ACC: Aortic cross-clamp

AKI: Acute kidney injury

ALAT: Alanine aminotransferase

ASAT: Aspartate aminotransferase

AVR: Aortic valve replacement

BUN: Blood urea nitrogen

CABG: Coronary artery bypass graft

COPD: Chronic obstructive pulmonary disease

CPB: Cardiopulmonary bypass

CRP: C-reactive protein

CVP: Central venous pressure

EF: Ejection fraction

EuroSCORE: European System of Cardiac Operative Risk Evaluation

FIO<sub>2</sub>: Fraction of inhaled oxygen

GFR: Glomerular filtration rate

GGT: Gamma-glutamyl transferase

IABP: Intracortical balloon pump

ICU: Intensive Care Unit

INR: Internationally normalized ratio

IQR: Interquartile range

IVC: Inferior vena cava

KDIGO: Kidney Disease: Improving Global Outcomes

LA: Left atrium

LV: Left ventricle

LVEDD: Left ventricle end-diastolic diameter

LVESD: Left-ventricle end-systolic diameter

MELD: Model for End-stage Liver Disease

MVR: Mitral valve replacement

NO: Nitric oxide

NYHA: New York Heart Association System

OPCAB: Off-pump coronary artery bypass  
PEEP: Positive end-expiratory pressure  
POCUS: Point-of-care ultrasound  
POP: Postoperative period  
RA: Right atrium  
RASA: Right atrium systolic area  
RBC: Red blood cell  
RCT: Randomized controlled trial  
ROC: Receiver operating characteristic  
ROS: Reactive oxygen species  
RR: Respiratory rate  
RV: Right ventricle  
STROBE: Strengthening the Reporting of Observational studies in Epidemiology  
TAPSE: Tricuspid Annular Plane Systolic Excursion  
TEE: Transesophageal echocardiography  
TTE: Transthoracic echocardiography  
TV: Tidal volume  
VExUS: Venous excess ultrasound  
VIS: Vasoactive-inotropic score  
VTI: Velocity-time integral  
v max: Maximal velocity

### 3. Introduction

#### 1. Fluid balance and fluid overload in critically ill patients

The accurate estimate of hemodynamic conditions is crucial in guiding the clinical management of the surgical patient in the postoperative period or during the long-term intensive care unit stay. (1) New protocols and guidelines put more emphasis on bolus and restrictive continuous fluid administration. This approach is ideal for critically ill patients as well, in whom increased endothelial permeability makes this approach more relevant. Malbrain uses the “R.O.S.E. conceptual model (Resuscitation, Optimization, Stabilization, Evacuation)”, which summarizes a dynamic approach to fluid therapy, maximizing benefits and minimizing harms. (2)

Progressing heart failure comes with endothelial dysfunction. According to recent studies, it may result from increased oxidative stress and activation of the adrenergic and the renin-angiotensin systems and from the production of inflammatory cytokines, which contribute to reduced bioavailability of nitric oxide (NO). Oxidative stress, determined by excess production of reactive oxygen species and impairment in the antioxidant defense, is responsible for both the decline of diffusible NO and the decrease in the concentration of essential co-factors of NO synthases. This leads to the compromised NO bioavailability and reduced vasodilatation. This leads to endothelial dysfunction. (3) (4)

Personalized fluid management is particularly important in cardiac surgery patients, especially those with heart failure, prolonged operative and aortic cross-clamp (ACC) time, or renal, pulmonary, or hepatic insufficiency.

A large amount of data in the international literature indicates that postoperative fluid overload in postoperative cardiac intensive care is associated with prolonged mechanical ventilation, higher vasoactive, inotropic support, and mortality. (5) (6) (7)

Fluid overload recognition and assessment require an accurate documentation of intakes and outputs; yet there is a wide difference in how it is evaluated. (8) There are several methods to estimate fluid status; however, most of the tests currently used are inaccurate. Heart failure is one of the main cardiac diseases leading to intensive care unit (ICU) admission (18.6%). (9) Organ congestion is a classical feature of heart failure that is caused by abnormal fluid redistribution into the extravascular space. In addition, during and after cardiac surgery the infusion of considerable amounts of intravenous fluids may be necessary due to cardiopulmonary bypass (CPB) induced inflammation, blood loss, myocardial depression, rhythm disturbances, and impaired vascular tone. Therefore, additional iatrogenic fluid overload is more common at ICU admission of these patients. Furthermore, patients with heart failure often suffer from extravascular over-hydration due to fluid re-distribution during intravascular fluid underload. This may lead to the administration of more fluid for resuscitation purposes, and this leads to more severe organ dysfunction (heart, lung, and kidneys), leading to a vicious circle of organ failure. Among calculating fluid balance (input and output) and checking hemodynamic markers (blood pressure, central venous pressure) protocols using ultrasound examinations are gaining more attention.

Inferior vena cava (IVC) measurements are commonly used by intensive care and emergency practitioners at point-of-care ultrasound (POCUS) examinations to predict fluid status. (10) (11-15) Using IVC measurements only can lead to inaccurate results, as they do not reflect the preload status of the patient's left ventricle accurately. The IVC can also be dilated in conditions such as tricuspid or mitral regurgitation, increased intrathoracic pressure with mechanical ventilation, increased right atrial pressure in pulmonary embolism or pulmonary hypertension, and in increased intra-abdominal pressure. Also, it shows a wide range of individual differences, especially in athletes. (16) Several recent protocols, such as the venous excess ultrasound examination (VExUS), use hepatic, portal, and renal measurements for Doppler ultrasound analysis. (17) The renal venous Doppler pattern is seen as a continuous monophasic flow. As venous congestion increases, there is a decrease of the systolic component of the wave with progression to a biphasic pattern and later a complete absence of systolic flow can be seen. The portal venous flow is normally monophasic with little variation. As venous

congestion increases, increasing amounts of pulsatility can be detected in the pattern. (18-20)

Several data in the literature suggest that abnormal hepatic, renal, and portal flows are good indicators of abnormal fluid shifts. Bhardwaj et al. evaluated their modified VExUS protocol in predicting AKI in patients with cardiorenal syndrome. They found a strong correlation between grades of VExUS and the stages of AKI. (11, 21)




Figure 1 A and B: Connection between the right ventricle, atrium, and the liver in case of right heart failure (own recording and figure) (5, 22)

## 2. Hepatic venous flow and abdominal congestion

Abdominal complications are well-known phenomena among cardiac surgery patients. Dong et al. observed over 2000 cardiac surgery cases, out of which 33 (1.4%) developed abdominal complications postoperatively, including cases of paralytic ileus, gastrointestinal hemorrhage, gastroduodenal ulcer perforation, acute calculus cholecystitis, hepatic dysfunction, and ischemic bowel diseases. (23)

The latest approaches are focusing on abdominal congestion postoperatively. Abnormal flow patterns in abdominal organs are correlated with elevated right-atrial pressure, thus providing additive information about end-organ congestion. In brief, congestion indicates poor outcomes in heart failure patients; therefore, a comprehensive assessment of venous congestion has prognostic significance. (24)

The relationship between the worse outcomes of cardiac diseases and liver dysfunction has been a well-recognized and described phenomenon for several decades. (25, 26) "Candesartan in Heart Failure: Assessment of Reduction in Mortality and Morbidity (CHARM)" showed in 2679 heart patients that abnormal total bilirubin, direct bilirubin, alkaline phosphatase, and albumin levels were predictive factors of the NYHA class (New York According to the Heart Association), the reduction of the ejection fraction, and the development of diabetes mellitus. (27)

The severity of liver congestion alone affects mortality in heart patients. Destruction of the zona pellucida caused by stasis leads to a fistula between the choledochal ducts and the sinusoids. Clinical symptoms can be poor, and in advanced liver failure we observe reduced coagulation factor and albumin synthesis. (28) (29)

In perioperative situations, many factors can influence hypoperfusion, hypoxia, or stagnation in the liver. The most important of these are the length of cardiopulmonary bypass, post-bypass low stroke volume, as well as the use of vasoactive, inotropic, immunosuppressive drugs, and blood products. Influencing factors that cannot be ruled out are mechanical circulatory support devices and the degree of resistance of the right ventricle and pulmonary vessels. It is also a known phenomenon that liver cirrhosis affects the hemodynamic parameters and leads to arterial vasodilatation. In addition, it also changes the fluid and electrolyte balance, which has a negative effect on the physiological regulation of circulation. (30)

We examined the hepatic venous Doppler ultrasound profile closely. There are three main hepatic veins—left, middle, and right—that separate the liver along craniocaudal planes. In 50% of cases, a well-defined right inferior hepatic vein accompanies the right hepatic vein. In about 70% of people, the left and middle hepatic veins merge to form a single vein before entering the IVC. We examined this common vein, right prior to its inflow into the IVC. The normal hepatic vein waveform has four components: a retrograde A

wave, an antegrade S wave, a transient V wave (which can be antegrade, retrograde, or neutral, but mainly retrograde), and an antegrade D wave.

The A wave corresponds to atrial contraction. With the tricuspid valve open, blood is pushed in two directions: antegrade toward the right ventricle and retrograde toward the IVC and into the hepatic veins. This results in a retrograde wave measured over the hepatic veins. As ventricular systole commences, the tricuspid valve closes and the retrograde velocity toward the hepatic veins begins to decrease and approach the baseline. During ventricular systole, not only the ventricular walls contract to push blood into the outflow tract, but there is also a movement of the tricuspid valve annulus toward the cardiac apex. These actions create a relative negative pressure in the atrium, causing antegrade blood flow out of the liver and into the heart during the S wave. In the normal heart, the largest amount of antegrade blood flow is during this phase. The V wave corresponds to atrial overfilling. As the ventricular contraction becomes less intense and the closed tricuspid valve begins to return to its original position, the atrium fills and blood flow velocity toward the heart decreases. The D wave begins as the tricuspid valve opens. During cardiac diastole, the right atrium and ventricle fill passively, with the antegrade flow of blood from the liver into the heart. In the normal patient, the velocity of this passive flow is almost always lower in magnitude than the velocity during the S wave.

There are a few standard differential diagnostic questions we must ask to analyze hepatic venous flow. We used the method applied by Scheinfeld et al. (31)



The first question we need to examine on the flow profile is whether we see any flow at all, and if so, whether it is in the right direction. If there is no flow or only retrograde flow can be seen, hepatic vein obstruction, such as Budd-Chiari syndrome, can be clearly concluded. Turbulent flow can be seen if the obstruction is not complete. If the flow is correct, its regularity must be checked. In the case of monophasic waves or a completely suppressed pattern, we should think primarily of liver tissue damage. This could be fatty liver, cirrhosis, or malignancy. If healthy phaticity is observed, the last step is to compare the magnitude of the D and S waves.

Normally, the S wave is greater than or equal to the D. This is because the strong systolic movement of the tricuspid annulus toward the apex of the heart causes a large anterograde flow toward the heart. We can see a deviation from this in cardiac pathologies when the

right side of the heart does not function properly, such as in tricuspid regurgitation, and in this case the diastolic anterograde flow will be greater than the systolic one.

If we take a closer look at the pathophysiology of these conditions, we can see that during atrial systole, the atrium contracts and simultaneously pumps blood into the ventricle and back toward the liver, and this can be seen as a small retrograde A wave on the Doppler ultrasound image of the hepatic veins. During ventricular systole, the ventricle contracts and the tricuspid annulus moves towards the apex of the heart, as has been mentioned above. However, because tricuspid regurgitation exists, blood flows retrograde into the atrium, vena cava, and liver following the flow path. This causes a decrease in the S wave or can even turn it retrograde in the liver. When the ventricles relax and the annulus returns to its original position, blood flows from the atrium again in a retrograde direction towards the liver, and this results in a retrograde V wave. Finally, during diastole, when the myocardium is relaxed and the tricuspid valve is open, blood passively flows from the liver into the right atrium, and this appears as a large, only anterograde D wave. (32) The picture of right heart failure is similar to tricuspid regurgitation. A slightly enlarged A wave may appear because of the increased backflow, since the abnormal right ventricle cannot adapt to the full atrial anterograde flow. These abnormalities have been observed since the 1980s. (33) (34)

We can divide the waves into retrograde (A and V) and anterograde (S and D) subgroups. If we observe these waves regarding the fluid balance, we can see in the normal hepatic venous blood flow waveform that the amplitude of the systolic component is higher than that of the diastolic component. For mild venous congestion, the amplitude of the systolic component is lower than that of the diastolic component. In severe congestion, there is a retrograde systolic component (opposite to the diastolic component). (35) (36, 37)






Figure 2 A and B: Hepatic venous flow (own recording and figure) (5, 22)

### 3. Postoperative right ventricle dysfunction, congestion, and acute kidney injury after cardiac surgery

Assessment of right ventricular (RV) function is crucial since RV failure with a reduced cardiac output (CO) is associated with compromised outcome in cardiac surgery. (38)

Right Ventricular (RV) output mostly derives from longitudinal shortening in normal hearts. However, following even uncomplicated cardiac surgery with preserved RV function, a significant and sustained decrease in longitudinal contraction has been observed. (39) RV dysfunction leads to systemic congestion.

Recent studies have recognized that increased renal backpressure caused by venous congestion is the major pathophysiological mechanism of renal dysfunction in postoperative cardiorenal syndrome. The systemic congestion generates increased renal venous pressure and reduces the arteriovenous gradient over renal circulation and thereby impairing renal blood flow. Moreover, independent from renal blood flow, increased renal venous pressure leads to renal parenchymal congestion, resulting in increased

interstitial pressure, which may compress all capillary and renal tubules, thereby reducing GFR in a physical way. (40) Acute kidney injury is a clinical syndrome manifested by rapid decline in kidney function and subsequent dysregulation of body electrolytes and volume, as well as abnormal retention of nitrogenous waste. The widely accepted Kidney Disease: Improving Global Outcome (KDIGO) definition of AKI is based on the change of serum creatinine and urine output: (41)

By the definition using creatinine stage 1 AKI means 1.5-1.9 times the baseline value or more than 0.3 mg/dL creatinine increment, stage 2 AKI 2-2.9 times, stage 3 AKI is more than 3 times increment or more than 4 mg/dL value. By urine output stage 1 means less than 0.5 ml/kg/h for 6-12 hours, over 12 hours is stage 2, and stage 3 means less than 0.3 ml/kg/h over 24 hours or 12 h anuria. (42)

Perioperative AKI is common and is associated with serious morbidity and mortality after cardiac surgery. The pathophysiology behind it is multifactorial. These include genetic predisposition, nephrotoxins, cardiopulmonary bypass (CPB) induced hemolysis, ischemic-reperfusion injury, complexity of cardiac surgery, oxidative stress, and inflammation. (43)

Examining the preoperative factors, advancing age is associated with decrease in renal function and estimated GFR, compromising normal renal physiology and increasing inclination to postoperative AKI. Furthermore, studies have found that women are more prone to develop AKI than men. (44)

Patients with comorbidities such as chronic obstructive pulmonary disease (COPD), diabetes mellitus, congestive heart failure, and chronic kidney disease (CKD) are prone to develop kidney injury following cardiac surgery. This may be due to impaired renal perfusion, damage to the endothelium, and reduced renal reserve. Moreover, these patients with these comorbidities are often given a variety of nephrotoxic drugs, such as nonsteroidal anti-inflammatory drugs (NSAIDs), angiotensin-converting enzyme inhibitors (ACEIs), and angiotensin receptor blockers (ARBs), which make these situations more severe.

In the intraoperative period, another potential nephrotoxin is free hemoglobin from CPB-induced hemolysis. CPB hemolyzes red blood cells and leads to the production of intravascular free hemoglobin, which depletes circulating haptoglobin and directly damages the renal endothelium. (45) During cardiac surgery, low cardiac output, potential hypothermia, possible bleeding complications, and the non-pulsatile flow of the CPB all cause reduced renal perfusion and can lead to AKI. The other fundamental cause of low renal perfusion is vasoconstriction. The endogenous way is the activation of the sympathetic nervous system and the renin-angiotensin system. The iatrogenic way is the use of vasopressin and catecholamines. Postoperatively, predominant factors influencing the development of AKI include hemodynamic instability, nephrotoxic, inotropic, and vasoconstrictor drugs, as well as systemic inflammation. Conventional on-pump CABG induces a systemic inflammatory response and is associated with elevated postoperative plasma concentrations of proinflammatory cytokines. A systematic review by Cheungpasitporn et al. (46), which included 33 randomized controlled trials (RCTs) showed a protective effect of OPCAB in reducing the risk of AKI, but the finding was challenged by the largest multicenter RCT by Lamy et al., (47) which found no significant difference between the OPCAB and on-pump CABG subgroups. In a prospective study, Bruins et al. reported a biphasic complement activation in cardiac surgery patients, which was activated not only during CPB but also during the first five days thereafter, which increased C-reactive protein levels, contributing to the second activation. (48) Other stimuli, such as endotoxemia, tissue injury, surgical trauma, and even left ventricular dysfunction, may play a role in the activation of the immune system. Inflammatory cytokines are formed, leading to renal tubular injury, which is getting worse by the activation of neutrophils and macrophages and the migration of lymphocytes into the renal parenchyma. Ischemia-reperfusion injury occurs and induces the production of reactive oxygen species (ROS). ROS activates the upregulation of pro-inflammatory transcription factors. (49)

AKI has been associated with an increased risk of sepsis, anemia, coagulopathy, and thus prolonged mechanical ventilation, ICU and hospital stay. (50) (51) The meta-analysis by Yu et al. concludes that the incidence of AKI after cardiac surgery is as high as 40%, and approximately 3% of patients require at least temporary renal replacement therapy. Patients with severe AKI are confronted with a three- to eightfold higher perioperative

mortality. (52) Messmer et al. showed in a meta-analysis that fluid overload and cumulative fluid balance were both associated with pooled mortality in a general postoperative population with AKI and non-AKI subgroups after surgery. (53) Malbrain analyzed 19,902 critically ill patients, and the cumulative fluid balance after one week of ICU stay was 4.4 L more positive in non-survivors compared to survivors. A restrictive fluid management strategy resulted in a less positive cumulative fluid balance of 5.6 L compared to controls after one week of ICU stay. (54) A restrictive fluid management was associated with a lower mortality compared to patients treated with a more liberal fluid management strategy. (2) (55) (56)

Petty showed an association between the development of AKI and a decrease in hepatic flow ratios on postoperative day 1, driven by low S-wave and high D-wave velocities. The presence of venous congestion was reflected by significantly elevated CVP values, which were independently associated with AKI on day 1. (57)

However, the influence of postoperative interventions, such as fluid therapy and the amount of the vasoactive drugs used has not been analyzed in relation to venous return. Additionally, there is no available data which calculates the retrograde and anterograde flow in the hepatic veins.

#### 4. Objectives

The aim of our studies was to investigate the perioperative factors that may be associated with increased hepatic venous congestion in the preoperative and postoperative periods in patients undergoing cardiac surgery.

The first objective was to investigate relationships between preoperative Doppler hepatic waveforms and echocardiographic parameters, ventilator settings, vasoactive medications, as well as laboratory parameters of renal and liver functions, especially focusing on renal parameters and AKI.

The second objective was to analyze the anterograde and retrograde flow of the hepatic vein. Our hypothesis was that the retrograde/anterograde ratio before and after cardiac surgery could both be helpful markers in the prediction of postoperative morbidity such as fluid overload, elevated vasoactive and inotrope need, worse left and right ventricle function, and more invasive ventilator settings.

## 5. Methods

### 1. Structure of the studies

All our studies were reported according to the STROBE statement. Both our main studies had approval from the Institutional Review Board of Semmelweis University (IRB 141/2018) and are registered with the ClinicalTrials.gov number NCT02893657. In the first prospective, observational study (study A), we enrolled 98 patients who underwent cardiac surgery between January 2018 and December 2019 in the Városmajor Heart and Vascular Center. In the second study (study B) 41 patients undergoing cardiac surgery between January 2021 and March 2021 were enrolled from the same tertiary heart center. Patients who had preoperative chronic kidney disease (defined as GFR under 30 mL/min/1.73 m<sup>2</sup>), hepatic cirrhosis, or portal vein thrombosis were excluded. Each patient who agreed to the participation signed the informed consent before the first echocardiographic investigation, which was usually 24 to 72 hours before the planned surgery.

Demographic data, perioperative echocardiographic parameters (preoperative and postoperative first day), and intraoperative variables (procedure type, cardiopulmonary bypass time) were collected. Postoperative factors (fluid balance, vasoactive requirement, ventilation time and parameters—respiratory rate (RR), tidal volume (TV), positive end-expiratory pressure (PEEP), fraction of inhaled oxygen (FIO<sub>2</sub>), and perioperative laboratory parameters (liver function, kidney function, albumin) were also used in the analysis. Laboratory results were collected during the preoperative period and on the first, second, and third postoperative days. We collected information on the predictors of the European System for Cardiac Operative Risk Evaluation (EuroSCORE II) and the Model for End-stage Liver Disease (MELD) scores, and the vasoactive-inotrope score (VIS). EuroSCORE II is a widely used risk stratification system for the cardiac surgical population using patient (clinical preoperative state, mobility), operation (urgency, operation on the thoracic aorta), and cardiac risk factors (LV function, recent myocardial infarction, pulmonary hypertension). (58) (59) VIS is calculated using vasopressor

(norepinephrine, epinephrine, vasopressin) and inotropic (dobutamine, dopamine, levosimendan, milrinone) medication doses. VIS was calculated using the highest doses of vasoactive and inotropic medications administered during the first 72 hours post-surgery. The MELD score has been used in transplant surgery since 2003 to estimate the need and urgency of liver transplants, which contains INR, creatinine, and bilirubin levels. The modified MELD-score (modMELD) differs from the standard MELD-score in that it considers the albumin level instead of the INR component, prioritizing the synthetic activity of the liver. Among cardiac and circulatory patients, many are under oral anticoagulation therapy, which greatly affects the INR and distorts the result, so there is an INR-excluding MELD-score too. An alternative MELD scoring system, MELD-Na may improve prognostic efficacy by incorporating low sodium levels, a commonly cited marker of increased mortality in patients with heart failure. (60, 61) We have calculated the AKI score by Thakar et al. – also containing risk factors in terms of patient (age, gender, diabetes, chronic obstructive pulmonary disease) and operation (procedure type, urgency) and preoperative creatinine level. (62) Subsequently, these scores were estimated and adjusted in our multivariable analyses.

$$\begin{aligned} \text{inotropic score (IS)} &= \text{dopamine dose (ug/kg/min)} + \text{dobutamine dose} \\ &\quad \text{(ug/kg/min)} + 100 \times \text{epinephrine dose (ug/kg/min)} \\ \text{vasoactive-inotropic score (VIS)} &= \text{IS} + 10 \times \text{milrinone dose (ug/kg/min)} + \\ &\quad 100 \times \text{norepinephrine dose (ug/kg/min)} + 10000 \times \text{vasopressine dose} \\ &\quad \quad \quad \text{(U/kg/min)} \end{aligned}$$

Figure 3: Calculation of the VIS score.

$$\begin{aligned} \text{MELD} &= 10 \times (1.12 \times \ln \text{INR} + 0.378 \times \ln \text{T. bili} + 0.957 \times \ln \text{Creat.} + 0.643) \\ \text{modMELD} &= 10 \times (1.12 \times \ln (1 + (4.1 - \text{albumin})) + 0.378 \times \ln \text{T. bili} + 0.957 \times \ln \\ &\quad \text{Creat.} + 0.643) \\ \text{MELD Na} &= \text{MELD} - \text{Na} - (0.025 \times \text{MELD} \times (140 - \text{Na})) + 140 \\ \text{MELD XINR} &= 5.1 \times \ln \text{T. bili} + 11.76 \times \ln \text{Creat.} + 9.44 \end{aligned}$$



Figure 4: Calculation of MELD-scores

Risk factor	Points
Female gender	1
Congestive heart failure	1
Ejection fraction less than 35%	1
Preoperative use of IABP	2
COPD	1
Insulin-requiring diabetes	1
Previous cardiac surgery	1
Emergency surgery	2
Valve surgery only	1
CABG+valve	2
Other cardiac surgery	2
Preoperative creatinine 1.2 to 2.1 mg/dl	2
Preoperative creatinine more than 2.1 mg/dl	5

Figure 5: AKI score by Thakar (62)

## 2. Ultrasound examinations

We recorded the maximum velocity of the four waves (A, S, V, D) and velocity-time integral (VTI). The ratio of maximum to retrograde compared to anterograde velocity and the ratio of retrograde VTI compared to anterograde VTIs were calculated. All recordings were made at the influx point of the hepatic veins into the inferior caval vein, and under the patients' inspiration.



Figures 6 A and B: Postoperative retrograde flow increment (own recording) (5, 22)

The echocardiographic investigations were performed by two cardiologists board-certified in echocardiography and were recorded on the same machine and analyzed by the same person. (Phillips CX50 – SG61102002). Standard 2D parameters were ejection fraction, tricuspid annular plane systolic excursion (TAPSE), the diameters of the atria and ventricles, and the occurrence of valve insufficiency.

The physicians in the postoperative period were blinded to the results of the hepatic flow measurements.

### 3. Main goals

In “study A” (Connection between preoperative hepatic venous profile and postoperative acute kidney injury) our primary outcome measured was postoperative AKI, as defined by the KDIGO criteria. In case of low population of AKI cases, we have focused on postoperative creatinine elevation (even in cases that did not reach the KDIGO criteria). We calculated the percentage change of the highest postoperative serum creatinine from the baseline preoperative concentration ( $\% \Delta Cr$ ) on the first three postoperative days. Baseline creatinine was defined as the preoperative creatinine level measured after the indexed hospital admission but before the surgery (usually 24 hours before the surgery). In “study B” (Connection between hepatic venous congestion and morbidity) our goal was to determine which factors are associated with the increased congestion of the liver as measured by Doppler ultrasound in patients undergoing cardiac surgery. Hepatic venous congestion was based on the changes of the ratios of the retrograde hepatic venous flow between the preoperative and the postoperative period, congestion meant elevated retrograde ratio.

### 4. Statistical analysis

Normality was tested with the Kolmogorov–Smirnov test. Skewed distributions are described as medians and interquartile ranges (interquartile range 25–75) and were compared using the Mann–Whitney U test. Continuous variables were first expanded with restricted cubic splines and were only used in linear form if the deviation from linearity was not significant, as indicated by the global F test ( $p > 0.05$ ). Multivariable models

were tested for multicollinearity. When we used binary classifier system (for example, VTI-growth and non-VTI growth subgroups) we tested the diagnostic ability with receiver operating characteristic curves. The effect of hepatic waves on the peak creatinine fractional change ( $\% \Delta \text{Cr}$ ) was evaluated using multivariable linear regression analysis and adjusted for patient- and surgery-related characteristics that included adjustment for significant perioperative variables. The variables selection was based on a set of previously published studies with a focus on kidney injury after cardiac surgery. The multivariable models were tested for multicollinearity (by volume inflation factors) between these clinical variables. The adjusted changes in  $R^2$  were reported after each step of the regression model to determine the contribution of each variable that was added to the model. (5, 22) Our power analysis indicated that with eight potential predictors of acute kidney injury, we needed to perform a prediction model with a minimum of 90 patients to avoid overfitting. In the second study our power analysis suggested 50 patients would be ideal.

Statistical tests were 2-sided, and  $p < 0.05$  was considered statistically significant. All statistical analyses were performed with SPSS software, Version 27.0 (IBM, Armonk, NY, USA).

## 6. Results

### 1. Connection between preoperative hepatic venous profile and postoperative acute kidney injury (study A)

#### 1. Descriptive statistics

Of the 98 patients, 66 (67%) were males, 32 (33%) were females. The median age of the patients was 69.8 years (interquartile range [IQR 25–75] 13 years). The most common type of cardiac surgery was coronary artery bypass grafting (CABG) surgery (40%) followed by aortic valve repair (AVR – 29%). Seventeen patients (17.3%) developed AKI according to the KDIGO criteria.

The average change in postoperative creatinine levels compared to baseline was an 8.7% increase from 83.2  $\mu\text{mol/L}$  (IQR 25–75: 7.8) to 91.9  $\mu\text{mol}$  (IQR 25–75: 12.8). Among these patents with AKI, there was a 72.1% increase from 78.4  $\mu\text{mol/L}$  (IQR 25–75: 8.7) to 137.0  $\mu\text{mol/L}$  (IQR 25–75: 32.7), and the ratio of postoperative/preoperative creatinine was 1.65 (1.51–1.81). We can see a significant difference between the AKI and non-AKI subgroups with Mann-Whitney U test ( $p=0.001$ ). There was a significant difference between the two subgroups in the age of the patients ( $p=0.015$ ).

Table 1: Basic demographic and preoperative clinical parameters

	<b>All patients (N=98)</b>	<b>AKI (n=17)</b>	<b>non-AKI (n=81)</b>	<b><i>p</i></b>
<b>Male</b>	66 (67%)	11 (62%)	55 (70%)	0.43
<b>Female (Nr)</b>	32 (33%)	6 (38%)	26 (30%)	0.16
<b>Age (years)</b>	68.8 (11.2)	69.1 (7.4)	63.5 (13.9)	0.015
<b>Diabetes</b>	20 (20%)	5 (25%)	15 (19%)	0.09
<b>NYHA III/IV</b>	41 (41%)	10 (58.8%)	31 (39%)	0.12
<b>EuroSCORE</b>	1.6 (0.9)	1.6 (1.0)	1.5 (0.7)	0.09
<b>Weight (kg)</b>	74.6 (8.1)	72.6 (7.1)	75.1 (9.12)	0.43
<b>Operation time (min)</b>	182.4 (39.1)	178.1 (41.1)	188.8 (39.1)	0.88
<b>Aorta cross clamp time (min)</b>	47.8 (7.1)	40.8 (9.1)	48.1 (7.6)	0.73
<b>Type of surgery</b>				
<b>AVR</b>	28 (29%)	6 (35%)	22 (26%)	0.11
<b>CABG</b>	39 (40%)	7 (41%)	32 (39%)	0.06
<b>MVR</b>	20 (20%)	3 (17%)	17 (20%)	0.09
<b>Combined</b>	12 (12%)	1 (7%)	11 (15%)	0.12
<b>Hemoglobin (g/l)</b>	137.1 (18.5)	135.3 (19.7)	138.4 (18.7)	0.61
<b>Albumin (g/l)</b>	41.4 (8.6)	38.8 (8.1)	43.7 (7.9)	0.83
<b>Platelets (1000/l)</b>	217.5 (62.2)	218.6 (64.9)	211.4 (58.7)	0.75
<b>CRP (mg/l)</b>	4.3 (3.3)	4.2 (3.9)	4.5 (3.3)	0.53
<b>INR</b>	1.9 (10.5)	1.9 (0.3)	1.1 (0.2)	0.34
<b>ASAT (U/l)</b>	20.9 (10.4)	18.8 (6.3)	21.2 (13.2)	0.34
<b>ALAT (U/l)</b>	31.4 (21.3)	34.2 (18.9)	30.4 (27.3)	0.56
<b>Creatinine (µmol/l)</b>	87.8 (20.1)	109.8 (25.1)	78.4 (17.7)	0.001

<b>Urea Nitrogen (mmol/l)</b>	6.4 (2.1)	7.3 (1.8)	5.4 (3.2)	0.18
<b>GFR (ml/min/1.73m<sup>2</sup>)</b>	79.7 (15.4)	70.2 (15.6)	76.0 (16.6)	0.004
<b>Bilirubin (μmol/l)</b>	17.8 (43.3)	21.4 (7.4)	11.8 (6.5)	0.31
<b>MELD</b>	7.1 (19.0)	7.50 (4.6)	5.3 (1.2)	0.50
<b>NYHA</b>	2.1 (0.5)	2.1 (0.4)	2.2 (0.5)	0.10
<b>EuroSCORE</b>	1.6 (0.9)	1.6 (0.9)	1.5 (1.0)	0.09

AKI: Acute kidney injury, AVR: Aortic valve repair, MVR: Mitral valve repair, CABG: Coronary artery bypass grafting, CRP: C-reactive protein, EuroSCORE: European System for Cardiac Operative Risk Evaluation INR: Internationally normalized ratio, ASAT: Aspartate aminotransferase, ALAT: Alanine aminotransferase, GFR: Glomerular filtration rate, MELD: Model for end-stage liver disease NYHA: New York Heart Association

There was no difference between the AKI and non-AKI patients in their standard echocardiographic parameters before and after surgery.

Table 2: Echocardiographic parameters

		<b>All patients (N=98)</b>	<b>AKI (n=17)</b>	<b>non-AKI (n=81)</b>	<b><i>p</i></b>
<b>EF</b>	<b>preoperative</b>	54.9 (10.6)	54.0 (11.6)	55.8 (12.3)	0.49
	<b>postoperative</b>	57.3 (8.1)	55.6 (8.9)	58.4 (7.1)	0.77
<b>TAPSE (mm)</b>	<b>preoperative</b>	23.6 (8.0)	22.8 (8.8)	24.1 (8.9)	0.4
	<b>postoperative</b>	13.4 (8.1)	13.4 (7.6)	14.6 (8.6)	0.27
<b>LVEDD (mm)</b>	<b>preoperative</b>	50.3 (6.1)	50.5 (6.2)	49.6 (5.5)	0.73
	<b>postoperative</b>	46.0 (8.3)	44.6 (9.1)	46.3 (7.1)	0.97
<b>LVESD (mm)</b>	<b>preoperative</b>	35.8 (5.1)	36.1 (4.3)	34.3 (5.3)	0.38
	<b>postoperative</b>	34.8 (7.8)	36.1 (7.4)	33.0 (6.4)	0.92
<b>RV (mm)</b>	<b>preoperative</b>	30.8 (8.1)	29.2 (8.8)	32.7 (6.5)	0.14
	<b>postoperative</b>	33.2 (9.0)	32.3 (9.0)	33.7 (9.1)	0.96
<b>LA (mm)</b>	<b>preoperative</b>	43.1 (8.3)	41.8 (8.5)	43.4 (7.0)	0.44
	<b>postoperative</b>	43.3 (6.3)	43.7 (8.6)	43.2 (3.7)	0.31
<b>RA (mm)</b>	<b>preoperative</b>	40.5 (7.6)	38.0 (7.2)	41.9 (8.7)	0.66
	<b>postoperative</b>	43.2 (6.2)	44.1 (7.0)	43.2 (6.7)	0.54

EF: ejection fraction, TAPSE: tricuspid annular plane systolic excursion LVEDD: left ventricle end-diastolic diameter, LVESD: left ventricle end-systolic diameter, RV: right ventricle, LA: left atrium, RA: right atrium

## 2. Univariable and multivariable regression

We have observed a tendency between the severity of KDIGO stages and the VTI ratio ( $p=0.045$ ) on ROC analysis. Maximum velocity ( $v_{max}$ ) and velocity-time-integral (VTI) of A, S, V, and D waves were analyzed regarding creatinine change in univariable and multivariable regression models. We also calculated the sum of retrograde (retro VTI and retro  $v_{max}$ ), the sum of anterograde flow (antero VTI and antero  $v_{max}$ ) and the ratio of retrograde and anterograde waves.

The variables were adjusted for age, EuroSCORE (European System for Cardiac Operative Risk Evaluation), AKI (acute kidney injury by Thakar et al) score, operation time, fluid balance, and vasoactive inotropic score (VIS). With univariable linear regression, we calculated the correlation between the preoperative parameters and the changes in postoperative creatinine levels compared to baseline. In the multivariable regression model, VTI and Vmax of the A wave correlated independently with the increase in the creatinine level. The VTI of the retrograde waves (A + D) and the VTI of the retrograde/antegrade waves ratio were independently associated with the increase in creatinine levels.

Table 3: Correlation between hepatic waves and changes in creatinine levels

	Univariable linear regression				Multivariable regression			
	<b>B</b>	<b>95% CI</b>		<b>p</b>	<b>B</b>	<b>95% CI</b>		<b>p</b>
<b>A vmax (m/s)</b>	0.640	0.332	0.948	<0.001	0.714	0.437	0.991	<0.001
<b>A VTI (cm)</b>	0.035	0.021	0.050	<0.001	0.038	0.025	0.051	<0.001
<b>S vmax (m/s)</b>	0.049	-0.153	0.251	0.631				
<b>S VTI (cm)</b>	-0.002	-0.011	0.007	0.691				
<b>V vmax (m/s)</b>	0.075	-0.127	0.277	0.462				
<b>V VTI (cm)</b>	0.011	-0.003	0.025	0.127				
<b>D vmax (m/s)</b>	0.151	-0.163	0.466	0.342				
<b>D VTI (cm)</b>	0.010	-0.005	0.024	0.177				
<b>Antero vmax (m/s)</b>	0.062	-0.080	0.204	0.388				
<b>Retro vmax (m/s)</b>	0.168	-0.009	0.345	0.062				
<b>Ratio vmax</b>	0.091	-0.124	0.307	0.402				
<b>Antero VTI (cm)</b>	0.001	-0.006	0.009	0.690				
<b>Retro VTI (cm)</b>	0.017	0.006	0.027	0.002	0.018	0.008	0.027	<0.001
<b>Ratio VTI</b>	0.218	0.086	0.351	0.002	0.233	0.111	0.356	<0.001

Among the postoperative variables characterizing the complications, the amount of transfusion in red-blood cell units and need for furosemide in the first 72 h were associated with changes in postoperative creatinine levels ( $\Delta\text{Cr}$ ).

Table 4: Postoperative parameters and correlations with changes in creatinine levels

	<b>All patients (N=98)</b>	<b>AKI (n=17)</b>	<b>non-AKI (n=81)</b>	<i>p</i>	<b>B</b>	<b>95% CI</b>		<i>p</i>
<b>Mechanical ventilation (hours)</b>	46.1 (9.5)	47.1 (9.2)	46.9 (10.1)	0.076	0.026	-0.002	0.054	0.066
<b>Need for vasopressor/ inotrope</b>	29 (32%)	11 (64%)	18 (23%)	0.234	0.282	-0.271	0.836	0.314
<b>VIS</b>	4.2 (8.2)	4.9 (8.1)	3.6 (8.2)	0.122	0.008	0.000	0.017	0.057
<b>Need for dialysis</b>	3 (3.7%)	2 (2.4%)	1 (1.3%)	0.09	0.021	-0.012	0.022	0.065
<b>ICU days</b>	2.2 (0.9)	2.3 (1.0)	2.2 (0.9)	0.123	0.277	-0.223	0.766	0.344
<b>Hospital days</b>	12.8 (5.6)	14 (6.5)	12 (8.7)	0.245	0.123	-0.156	0.544	0.231
<b>Transfusion RBC (U)</b>	0 (0-2)	2.2 (3.1)	0.9 (1.9)	0.002	0.043	0.008	0.078	0.016
<b>CVP (max) (Hgmm)</b>	9.0 (2.5)	11.0 (2.0)	8.0 (1.2)	0.322	0.009	-0.005	0.023	0.197
<b>Fluid balance (ml/kg)</b>	34.8 (29.1)	50.1 (40.1)	19.6 (3.4)	0.098	0.001	-0.005	0.023	0.068
<b>Furosemide (mg)</b>	25.0 (10.0)	40.0 (10.0)	20.0 (10.0)	0.001	0.004	0.003	0.006	0.001

RBC: Red blood cell, ICU: intensive care unit, CVP: central venous pressure

## 2. Connection between hepatic venous congestion and morbidity (study B)

### 1. Descriptive statistics

Of the 41 patients, 20 (48.7%) were male. The median age of the patients was 65.9 years (IQR: 59.8–69.9 years). The average increase in the retrograde/antegrade wave VTI ratio was 0.04 (from preoperative  $15.5/20.8 = 0.77$  to postoperative  $19.7/24.4 = 0.81$ ). After examining the waves, we found that most of the population had an increased retrograde VTI growth (24 - 59%), compared to the baseline preoperative values. In the retrograde VTI growth subgroup, delta D decreased, and there was a tendency toward an increase in delta A and delta V waves. There was no significant difference between the two subgroups among the basic demographic and clinical parameters.


Table 5: Demographic and basic clinical parameters



	<b>All patients (N=41)</b>	<b>No retrograde VTI growth (n=17)</b>	<b>Retrograde VTI growth (n=24)</b>	<b><i>p</i></b>
<b>Age (years)</b>	65.9 (10.8)	63.7 (11.3)	67.1 (10.4)	0.272
<b>Weight (kg)</b>	72.0 (10.2)	73.4 (11.1)	71.1 (10.3)	0.845
<b>Diabetes</b>	19 (46.3%)	9 (53%)	10 (41.6%)	0.53
<b>Sex female</b>	21 (51.2%)	9 (53%)	12 (50%)	0.466
<b>EUROSCORE II</b>	4.7 (1.3)	4.9 (0.9)	4.5 (1.0)	0.197
<b>NYHA III/IV</b>	24 (58.5%)	10 (58.8%)	14 (58.3%)	0.456
<b>Operation time (min)</b>	182.4 (39.1)	178.1 (41.1)	188.8 (39.1)	0.88
<b>Aorta cross-clamp time (min)</b>	47.8 (7.1)	40.8 (9.1)	48.1 (7.6)	0.73
<b>Operation type</b>				
<b>MVR</b>	9 (22%)	3 (17.6%)	6 (25%)	0.234
<b>AVR</b>	14 (34.1%)	6 (35.2%)	8 (33.3%)	0.199
<b>CABG</b>	15 (36.6%)	7 (41.1%)	8 (33.3%)	0.342
<b>Combined</b>	3 (7.3%)	1 (5.9%)	2 (8.4%)	0.544

NYHA: New York Heart Association, MVR: Mitral valve repair, AVR: Aortic valve repair, CABG: Coronary artery bypass graft, EUROSCORE II: European System for Cardiac Operative Risk Evaluation II

Table 6: Changes in hepatic waves

	<b>non-VTI growth (n=17)</b>	<b>VTI growth (n=24)</b>	<b><i>p</i></b>
<b>Pre A (cm)</b>	9.1 (7.7)	8.8 (5.3)	0.434
<b>Pre S (cm)</b>	12.2 (9.9)	13.9 (9.9)	0.275
<b>Pre V (cm)</b>	6.4 (4.3)	5.3 (4.5)	0.209
<b>Pre D (cm)</b>	8.6 (4.2)	13.3 (6.7)	0.026
<b>Pre VTI ratio</b>	1.01 (0.02)	0.59 (0.01)	0.098
<b>Post A (cm)</b>	8.5 (5.7)	9.8 (5.6)	0.239
<b>Post S (cm)</b>	14.6 (9.0)	12.4 (8.1)	0.22
<b>Post V (cm)</b>	6.9 (4.1)	7.5 (6.6)	0.375
<b>Post D (cm)</b>	10.1 (7.3)	10.1 (6.9)	0.425
<b>Post VTI ratio</b>	0.65 (0.03)	0.91 (0.03)	0.087
<b>Delta A (cm)</b>	-0.64 (0.01)	0.99 (0.05)	0.08
<b>Delta S (cm)</b>	2.5 (1.1)	-1.4 (0.02)	0.08
<b>Delta V (cm)</b>	-0.01 (0.01)	2.28 (1.2)	0.117
<b>Delta D (cm)</b>	1.3 (0.2)	-2.2 (1.2)	0.007
<b>Delta VTI ratio</b>	-0.35 (0.01)	0.31 (0.02)	0.002





Figures 7 A, B and C: Changes between the waves in the preoperative and postoperative periods

Examining the laboratory parameters, the retrograde VTI ratio (elevated congestion) subgroup had significantly higher blood urea nitrogen and creatine levels.

Table 7: Laboratory parameters at preoperative and postoperative period's 24h.

	<b>non-VTI growth (n=17)</b>	<b>VTI growth (n=24)</b>	<i>p</i>	<b>non-VTI growth (n=17)</b>	<b>VTI growth (n=24)</b>	<i>p</i>
	<b>Preoperative</b>			<b>Postoperative</b>		
<b>Creatinine (mmol/l)</b>	83.2 (13.3)	85.5 (9.8)	0.381	78.1 (21.2)	93.5 (38.7)	0.07
<b>BUN (umol/l)</b>	5.9 (1.9)	6.3 (4.3)	0.235	4.75 (3.2)	6.4 (2.3)	0.02
<b>AST (U/l)</b>	22.1 (4.3)	34.1 (10.1)	0.135	55.6 (19.8)	47.5 (19.9)	0.297
<b>ALT (U/l)</b>	20.3 (11.1)	22.8 (4.9)	0.303	24.1 (10.1)	18.5 (8.9)	0.103
<b>Bilirubin (umol/l)</b>	11.04 (6.1)	11.07 (7.1)	0.462	11.3 (6.2)	11.1 (5.9)	0.368
<b>GGT (U/l)</b>	35.5 (11.1)	32.9 (13.2)	0.103	43.4 (9.1)	22.8 (8.9)	0.301
<b>GFR (ml/min/1.73 m2)</b>	71.2 (13.4)	65.3 (23.3)	0.112	74.4 (14.3)	55.1 (19.8)	0.001

After comparing the echocardiographic parameters between the two subgroups, the postoperative right ventricular and atrial diameters were significantly greater in the “retrograde VTI growth” group. The postoperative ejection fraction and the decrease in ejection fraction were significantly different between the two groups. ( $p = 0.001$  and  $0.009$ ).

Table 8: Changes in echocardiographic parameters

	<b>non-VTI growth (n=17)</b>	<b>VTI growth (n=24)</b>	<i>p</i>	<b>non-VTI growth (n=17)</b>	<b>VTI growth (n=24)</b>	<i>p</i>
	Preoperative			Postoperative		
<b>EF</b>	53.7 (10.5)	51.5 (12.4)	0.277	59.4 (11.2)	48.5 (10.7)	0.001
<b>TAPSE (mm)</b>	24.6 (5.1)	22.6 (5.9)	0.127	14.3 (4.2)	15.2 (5.4)	0.251
<b>LVEDD (mm)</b>	51.1 (9.2)	53.1 (10.4)	0.255	44.6 (4.1)	47.4 (6.8)	0.08
<b>LVESD (mm)</b>	39.9 (14.1)	38 (10.1)	0.431	28.4 (9.1)	33.3 (8.7)	0.022
<b>RV (mm)</b>	32.7 (4.4)	32.9 (3.9)	0.457	32.5 (5.1)	35.1 (4.9)	0.01
<b>LA1 (mm)</b>	44 (8.5)	46.6 (8.0)	0.168	43.1 (8.7)	45.2 (8.8)	0.06
<b>LA2 (mm)</b>	49 (10.2)	52.1 (10.1)	0.166	54.1 (8.7)	53.9 (9.1)	0.09
<b>RA1 (mm)</b>	42.7 (6.3)	42.4 (7.1)	0.451	39.5 (6.7)	43.9 (5.9)	0.01
<b>RA2 (mm)</b>	49.3 (9.1)	47.3 (7.9)	0.254	53.5 (6.9)	55.1 (7.1)	0.144
<b>RASA (mm<sup>2</sup>)</b>	2131.1 (644.1)	1916.2 (577.6)	0.134	2314.2 (498.2)	2278.5 (514.3)	0.646
<b>Delta EF</b>	2.8 (19.4)	-12.1 (3.2)	0.009			

## 2. Regressions

After analyzing the respiratory parameters and ventilation times, we found that the VTI growth subgroup had longer ventilation duration (the median value was over 24 h) and needed higher positive end-expiratory pressures ( $p = 0.003$ )

Table 9: Ventilation parameters

	<b>non-VTI growth (n=17)</b>	<b>VTI growth (n=24)</b>	<b><i>p</i></b>
<b>Mechanical ventilation over than 24 hours</b>	7 (41.1%)	13 (54.1%)	0.091
<b>Resp. time (hours)</b>	20.9 (2.1)	25 (3.2)	0.081
<b>Tidal volume (ml)</b>	470.5 (43.5)	490.5 (44.0)	0.079
<b>RR (/min)</b>	13.4 (2.0)	12.9 (1.9)	0.122
<b>PEEP (cmH2O)</b>	6.5 (1.3)	7.7 (1.9)	0.003
<b>Flo2 (%)</b>	38.5 (4.5)	39 (4.8)	0.051

In the univariable linear regression model, the postoperative/preoperative (baseline) delta VTI ratio of the hepatic venous ultrasound correlated with fluid balance, maximum central venous pressure, and delta ejection fraction.

Table 10: Regressions with postoperative/preoperative VTI ratio

	<b>B</b>	<b>95% CI</b>		<b><i>p</i></b>
<b>Delta EF</b>	-0.365	-0.022	-0.002	0.022
<b>Fluid balance/body weight at POP 24 h (ml/kg)</b>	0.341	0.001	0.021	0.022
<b>CVP POP 24 h (mmHg)</b>	0.546	0.052	0.213	0.002

## 7. Discussion

### 1. Perioperative fluid overload

We found that in the first postoperative 24 hours, increased hepatic venous retrograde flow was associated with worse ejection fraction, higher positive fluid balance, increased right atrial and ventricular diameter, and higher central venous preload pressure.

In the postoperative period, preload optimization can lead to fluid overload, particularly in low cardiac output states, preexisting or newly developed right-sided heart failure, and pulmonary hypertension. In the last decade, this connection between congestive heart failure, right ventricular failure, and liver dysfunction has been examined closer. Right ventricular failure can lead to liver dysfunction ranging from mild enzyme elevations to severe liver fibrosis. Congestion can impair renal function and cause gastrointestinal ischemia due to high backing pressure and a lower effective organ pressure gradient. Progressive increase in venous volume eventually causes the venous walls stretching to the maximum. At this point, the pressure transmission will be significantly increased, leading to peripheral venous congestion. (63) Increased right atrial pressure can be transmitted backwards across the venous tree and lead to congestive organ dysfunction. (64) This can manifest as congestive hepatopathy, delirium from congestive encephalopathy, and acute kidney injury from intracapsular oedema. (65) (66) (67)

Over the last 15 years, several dynamic tests have been developed to predict fluid status and fluid responsiveness in critically ill patients. These tests are based on the principle of inducing short-term changes in cardiac preload, using heart-lung interactions, the passive leg raise or by the infusion of small volumes of fluid, and to observe the resulting effect on cardiac output. Pulse pressure and stroke volume variations were first developed, but they are reliable only under strict conditions. (68-70) The variations in vena caval diameters share many limitations. The passive leg-raising test is now supported by solid evidence and is more frequently used. (71) (72, 73) More recently, the end-expiratory occlusion test was proposed, which can easily be performed in ventilated patients. Unlike the traditional fluid challenge, these dynamic tests do not lead to fluid overload. (74) (75)

Changes in renal venous flow patterns were associated with one-year mortality in patients with advanced heart failure independent from right atrial pressure. (67) The flow of the portal and hepatic veins was also examined. Among heart failure patients, those with reversed V waves had higher right atrial pressure and were at higher risk of adverse prognosis. Reversed V wave was also found to be an independent predictor of cardiac events. (76) Examining the portal vein pulsatility over 50% was an independent negative predictive value of worse outcomes among acute heart failure patients. (20)

These are new diagnostic concepts for practice in the early detection of fluid overload. It may be separate and not accompanying right-sided heart failure. In our study, we measured VTIs instead of the maximum velocity (v max) of the waveforms, which may be more useful for volume estimation. Using ratios to express retrograde and antero-grade flows has the advantage of removing the effects of vessel shape and diameter and reducing potential errors.

Our results suggest that an increase in retrograde flow was associated with the severity of fluid overload. It is not clear whether fluid overload leads to increased retrograde flow, or it is caused by pre-existing venous insufficiency (we can see this at lower extremity varicosities), or both.

## 2. Mechanical ventilation

Fast track management after cardiac surgery can be beneficial, as early return of spontaneous breathing and extubation promotes negative thoracic pressure, increasing the venous return, the preload of the heart, and helps maintaining cardiac output. (5) Negative pressure can also help reduce hepatic afterload, whereas fluid overload, prolonged ventilation, and edema formation worsen venous congestion. A special proof for this physiological phenomenon is patients with univentricular physiology (Fontane circulation), where early weaning from mechanical ventilation and return to spontaneous breathing is key to maintain cardiac output.

Early extubation is associated with lower cardiovascular morbidity, such as myocardial ischemia and lower post-surgical vasopressor requirement. (77) Prolonged positive pressure ventilation is associated with a decreased preload, reduction, and redistribution

of cardiac output, decreased splanchnic blood flow, and reduced blood supply to the liver and kidneys.

Our data suggests the increment of the retrograde hepatic venous flow is associated with longer ventilation duration (the median value was over 24 h) and a strong correlation with higher positive end-expiratory pressures.

### 3. Echocardiography

Echocardiography is a widely used and accepted technique in the perioperative management of patients undergoing cardiac surgery. In recent protocols echocardiography, with both first and second level parameters, offers the possibility to accurately analyze the right ventricle and optimize patient management. (78) (79) Echocardiography, both transthoracic (TTE) and transesophageal (TEE), may help to identify cardiac tamponade after surgery. (78) A meta-analysis by Heiberg (80) reported that echocardiography is feasible, whether performed by a novice or expert, and frequently resulted in important changes in the diagnosis of cardiac abnormalities and their management. Among the standard echocardiography, several point-of-care ultrasound markers (POCUS) have been identified recently as potential markers to detect venous congestion and fluid overload. (81) In our studies, we have used a combination of these measurements. By comparing the echocardiographic parameters between the venous congestion and non-congestion subgroups, we found significantly dilated right atrium and ventricle and lower ejection fraction in the assumed congestive subgroup.

### 4. Acute kidney injury

We found a significant correlation between the ratio of retrograde waves compared to anterograde hepatic venous waves, as well as VTI and maximum velocity of the A wave and postoperative AKI. The strong relationship of the retrograde A wave and the retrograde to anterograde ratio with AKI remained statistically significant after adjusting for patient- and procedure-related characteristics. VTI, but not maximum velocity, of the

retrograde and antegrade waves, showed a significant correlation with AKI. This also suggest that VTI is more effective in predicting volume dimensions than velocity. The severity of hepatic venous congestion was found to be a predictor of AKI in a prospective study among cardiac surgery patients. (82) (83)

Postoperative acute kidney injury is a challenge to clinicians at intensive care units. (49) (84) Various studies have measured the maximum velocity of the S and D waves and the severity of venous congestion by the S and D wave ratios preoperatively, during intensive care and after discharge. Based on our results, we found that the magnitude of the retrograde waves in hepatic venous circulation is not solely attributed to the severity of right heart failure. Among these parameters (tricuspid regurgitation, TAPSE, or right ventricular diameters) only the right atrial systolic area showed an association with the retrograde waves of the hepatic vein.

We chose AKI and an increase in creatinine levels as the main outcomes in our study. In the multivariable model, we also adjusted the retrograde waveforms for age, EuroSCORE II, GFR, and operative factors, and venous congestion remained significant. Our results indicate that the severity of venous congestion should be evaluated preoperatively, but it is also advised to follow it up in the postoperative period.

## 5. Other liver parameters

Severe congestive heart failure is associated with two distinct forms of liver dysfunction: passive congestion and acute hepatocellular necrosis that is caused by inadequate perfusion. Fibrosis may result from prolonged recurrent congestive heart failure. Ischemic hepatitis (shock liver) usually manifests as asymptomatic elevation of the serum aminotransferase (ASAT, ALAT) levels after an episode of hypotension, such as acute viral hepatitis. (85) Calculation of MELD and its modifications has been associated with increased mortality and morbidity in end-stage heart failure and in patients after heart transplantation. (86) (87-89) In our study population, there were no differences between MELD scores in AKI and non-AKI patients, but the association between bilirubin and retrograde A waves draws attention to possible hepatic synthetic dysfunction in cases of fluid overload and supranormal vasoactive drug administration.

## 8. Limitations

Our studies have limitations. First, it was a single center study. In the planning of the first study, we did not think about continuing it in the postoperative period. A couple of perioperative factors, such as positive fluid balance, vasopressor use, or positive intrathoracic pressure ventilation, can significantly influence the anterograde/retrograde ratios simultaneously. Therefore, we thought that the retrograde/anterograde ratio measurement would be the least biased choice. Furthermore, ultrasound methodology always has observational bias.

## 9. Conclusions

We can conclude that congestion and/or abdominal venous insufficiency in the liver can predict worse postoperative renal function. Abdominal ultrasound is an easily available and, most importantly, noninvasive tool, which is gaining more and more space in emergency and intensive care units. Monitoring the hepatic veins can easily be performed, together with standard preoperative echocardiography it may add more information about the liver and kidney, than laboratory parameters alone. Our new finding was that it is also advised to quantitatively express our measures and not only with flow pattern types.

Moreover, measurement of venous congestion by ultrasound seems to be an important tool in guiding personalized fluid management. Volume and pressure monitoring on the arterial and macrohemodynamic side does not provide sufficient hemodynamic data, and central venous pressure alone provides inadequate information about the venous side of the circulation and all available monitoring methods are extremely invasive or have serious limitations. Monitoring abdominal venous waveforms is an easy and inexpensive noninvasive method that can help to detect fluid overload at an early phase.

## 10. Summary

This thesis has set out to determine which factors are associated with the increased congestion of the liver, as measured by Doppler ultrasound in patients undergoing cardiac surgery, and the effect of this abdominal congestion on postoperative morbidity, especially on acute kidney injury.

- In total, we examined 139 patients who underwent an elective cardiac surgery. In study A, of the 98 patients, 66 (67%) were males, 32 (33%) were females. The median age of the patients was 69.8 years (interquartile range [IQR 25–75] 13 years). In study B, of the 41 patients, 20 (48.7%) were male. The median age of the patients was 65.9 years (IQR: 59.8–69.9 years).
- The postoperative/preoperative (baseline) delta VTI ratio of the hepatic venous ultrasound correlated with fluid balance, maximum central venous pressure, and delta ejection fraction.
- The postoperative retrograde/antegrade VTI ratio increment was associated with longer ventilation duration and needed higher positive end-expiratory pressures.
- The postoperative right ventricular and atrial diameters were significantly larger in the subgroup with greater retrograde/antegrade VTI ratio.
- In “study A” (Connection between preoperative hepatic venous profile and postoperative acute kidney injury) our primary outcome measured was postoperative AKI, as defined by the KDIGO criteria. Because of the low number of AKI cases, we have focused on postoperative creatinine elevation (even in cases that did not reach KDIGO criteria).
- VTI and Vmax of the A wave and the VTI of the retrograde waves (A + D), and the VTI of the retrograde/antegrade waves ratio correlated independently with the increase in the creatinine level.

Measurement of venous congestion by ultrasound seems to be an important tool in guiding personalized fluid management. Monitoring abdominal venous waveforms is an

easy and inexpensive noninvasive method that can help to detect fluid overload at an early phase.

## 11. References

1. De Backer D, Aissaoui N, Cecconi M, Chew MS, Denault A, Hajjar L, How can assessing hemodynamics help to assess volume status? *Intensive Care Med.* 2022. Epub 2022/08/10. doi: 10.1007/s00134-022-06808-9. PubMed PMID: 35945344; PubMed Central PMCID: PMC9363272.
2. Malbrain MLNG, Langer T, Annane D, Gattinoni L, Elbers P, Hahn RG, Intravenous fluid therapy in the perioperative and critical care setting: Executive summary of the International Fluid Academy (IFA). *Ann Intensive Care.* 2020;10(1):64. Epub 20200524. doi: 10.1186/s13613-020-00679-3. PubMed PMID: 32449147; PubMed Central PMCID: PMC7245999.
3. Parodi O, De Maria R, Roubina E. Redox state, oxidative stress and endothelial dysfunction in heart failure: the puzzle of nitrate-thiol interaction. *J Cardiovasc Med (Hagerstown).* 2007;8(10):765-74. Epub 2007/09/22. doi: 10.2459/JCM.0b013e32801194d4. PubMed PMID: 17885513.
4. Devaux C, Varin R, Mulder P, Richard V, Thuillez C. [Oxidative stress and endothelial dysfunction in heart failure]. *Therapie.* 2001;56(5):575-81. Epub 2002/01/25. PubMed PMID: 11806296.
5. Eke C, Szabó A, Nagy Á, Szécsi B, Szentgróti R, Dénes A, Association between Hepatic Venous Congestion and Adverse Outcomes after Cardiac Surgery. *Diagnostics (Basel).* 2022;12(12). Epub 20221215. doi: 10.3390/diagnostics12123175. PubMed PMID: 36553182; PubMed Central PMCID: PMC9777079.
6. Cecconi M, Hernandez G, Dunser M, Antonelli M, Baker T, Bakker J, Fluid administration for acute circulatory dysfunction using basic monitoring: narrative review and expert panel recommendations from an ESICM task force. *Intensive Care Med.* 2019;45(1):21-32. Epub 20181119. doi: 10.1007/s00134-018-5415-2. PubMed PMID: 30456467.
7. Seo Y, Nakatsukasa T, Sai S, Ishizu T, Iida N, Yamamoto M, Clinical implications of organ congestion in heart failure patients as assessed by ultrasonography. *Cardiovasc Diagn Ther.* 2018;8(1):57-69. Epub 2018/03/16. doi:

10.21037/cdt.2017.07.05. PubMed PMID: 29541611; PubMed Central PMCID: PMC5835652.

8. Claire-Del Granado R, Mehta RL. Fluid overload in the ICU: evaluation and management. *BMC Nephrol.* 2016;17(1):109. Epub 2016/08/04. doi: 10.1186/s12882-016-0323-6. PubMed PMID: 27484681; PubMed Central PMCID: PMC54970195.

9. Waskowski J, Michel MC, Steffen R, Messmer AS, Pfortmueller CA. Fluid overload and mortality in critically ill patients with severe heart failure and cardiogenic shock-An observational cohort study. *Front Med (Lausanne).* 2022;9:1040055. Epub 2022/12/06. doi: 10.3389/fmed.2022.1040055. PubMed PMID: 36465945; PubMed Central PMCID: PMC9712448.

10. Beaubien-Souligny W, Rola P, Haycock K, Bouchard J, Lamarche Y, Spiegel R, Quantifying systemic congestion with Point-Of-Care ultrasound: development of the venous excess ultrasound grading system. *Ultrasound J.* 2020;12(1):16. Epub 2020/04/10. doi: 10.1186/s13089-020-00163-w. PubMed PMID: 32270297; PubMed Central PMCID: PMC7142196.

11. Bhardwaj V, Vikneswaran G, Rola P, Raju S, Bhat RS, Jayakumar A, Combination of Inferior Vena Cava Diameter, Hepatic Venous Flow, and Portal Vein Pulsatility Index: Venous Excess Ultrasound Score (VEXUS Score) in Predicting Acute Kidney Injury in Patients with Cardiorenal Syndrome: A Prospective Cohort Study. *Indian J Crit Care Med.* 2020;24(9):783-9. Epub 2020/11/03. doi: 10.5005/jp-journals-10071-23570. PubMed PMID: 33132560; PubMed Central PMCID: PMC7584837.

12. Singh S, Koratala A. Utility of Doppler ultrasound derived hepatic and portal venous waveforms in the management of heart failure exacerbation. *Clin Case Rep.* 2020;8(8):1489-93. Epub 2020/05/20. doi: 10.1002/ccr3.2908. PubMed PMID: 32884781; PubMed Central PMCID: PMC7455438.

13. Singh Y, Tissot C, Fraga MV, Yousef N, Cortes RG, Lopez J, International evidence-based guidelines on Point of Care Ultrasound (POCUS) for critically ill neonates and children issued by the POCUS Working Group of the European Society of Paediatric and Neonatal Intensive Care (ESPNIC). *Crit Care.* 2020;24(1):65. Epub 2020/02/26. doi: 10.1186/s13054-020-2787-9. PubMed PMID: 32093763; PubMed Central PMCID: PMC7041196.

14. Sultan LR, Sehgal CM. A Review of Early Experience in Lung Ultrasound in the Diagnosis and Management of COVID-19. *Ultrasound Med Biol.* 2020;46(9):2530-45. Epub 20200525. doi: 10.1016/j.ultrasmedbio.2020.05.012. PubMed PMID: 32591166; PubMed Central PMCID: PMC7247506.
15. Zieleskiewicz L, Lopez A, Hraiech S, Baumstarck K, Pastene B, Di Bisceglie M, Bedside POCUS during ward emergencies is associated with improved diagnosis and outcome: an observational, prospective, controlled study. *Crit Care.* 2021;25(1):34. Epub 2021/01/24. doi: 10.1186/s13054-021-03466-z. PubMed PMID: 33482873; PubMed Central PMCID: PMC7825196.
16. Galindo P, Gasca C, Argaiz ER, Koratala A. Point of care venous Doppler ultrasound: Exploring the missing piece of bedside hemodynamic assessment. *World J Crit Care Med.* 2021;10(6):310-22. Epub 20211109. doi: 10.5492/wjccm.v10.i6.310. PubMed PMID: 34888157; PubMed Central PMCID: PMC78613717.
17. Argaiz ER. VExUS Nexus: Bedside Assessment of Venous Congestion. *Adv Chronic Kidney Dis.* 2021;28(3):252-61. Epub 2021/12/16. doi: 10.1053/j.ackd.2021.03.004. PubMed PMID: 34906310.
18. Koratala A. Venous congestion assessment using point-of-care Doppler ultrasound: Welcome to the future of volume status assessment. *Clin Case Rep.* 2021;9(3):1805-7. Epub 20210121. doi: 10.1002/ccr3.3840. PubMed PMID: 33768948; PubMed Central PMCID: PMC7981685.
19. Rola P, Miralles-Aguilar F, Argaiz E, Beaubien-Souligny W, Haycock K, Karimov T, Clinical applications of the venous excess ultrasound (VExUS) score: conceptual review and case series. *Ultrasound J.* 2021;13(1):32. Epub 2021/06/20. doi: 10.1186/s13089-021-00232-8. PubMed PMID: 34146184; PubMed Central PMCID: PMC8214649.
20. Torres-Arrese M, Mata-Martínez A, Luordo-Tedesco D, García-Casasola G, Alonso-González R, Montero-Hernández E, Usefulness of Systemic Venous Ultrasound Protocols in the Prognosis of Heart Failure Patients: Results from a Prospective Multicentric Study. *J Clin Med.* 2023;12(4). Epub 20230206. doi: 10.3390/jcm12041281. PubMed PMID: 36835816; PubMed Central PMCID: PMC9966251.

21. Gupta S, Tomar DS. VEXUS-The Third Eye for the Intensivist? *Indian J Crit Care Med.* 2020;24(9):746-7. Epub 2020/11/03. doi: 10.5005/jp-journals-10071-23582. PubMed PMID: 33132551; PubMed Central PMCID: PMCPMC7584833.
22. Eke C, Szabó A, Nagy Á, Párkányi B, Kertai MD, Fazekas L, Association between Preoperative Retrograde Hepatic Vein Flow and Acute Kidney Injury after Cardiac Surgery. *Diagnostics (Basel).* 2022;12(3). Epub 20220312. doi: 10.3390/diagnostics12030699. PubMed PMID: 35328250; PubMed Central PMCID: PMCPMC8946915.
23. Dong G, Liu C, Xu B, Jing H, Li D, Wu H. Postoperative abdominal complications after cardiopulmonary bypass. *J Cardiothorac Surg.* 2012;7:108. Epub 2012/10/11. doi: 10.1186/1749-8090-7-108. PubMed PMID: 23046511; PubMed Central PMCID: PMCPMC3493288.
24. D'Andrea A, Del Giudice C, Fabiani D, Caputo A, Sabatella F, Cante L, The Incremental Role of Multiorgan Point-of-Care Ultrasounds in the Emergency Setting. *Int J Environ Res Public Health.* 2023;20(3). Epub 2023/02/12. doi: 10.3390/ijerph20032088. PubMed PMID: 36767456; PubMed Central PMCID: PMCPMC9915087.
25. Auer J. What does the liver tell us about the failing heart? *European heart journal.* 2013;34(10):711-4. Epub 2012/12/19. doi: 10.1093/eurheartj/ehs440. PubMed PMID: 23257947.
26. Cogger VC, Fraser R, Le Couteur DG. Liver dysfunction and heart failure. *Am J Cardiol.* 2003;91(11):1399. doi: 10.1016/s0002-9149(03)00370-9. PubMed PMID: 12767448.
27. Allen LA, Felker GM, Pocock S, McMurray JJ, Pfeffer MA, Swedberg K, Liver function abnormalities and outcome in patients with chronic heart failure: data from the Candesartan in Heart Failure: Assessment of Reduction in Mortality and Morbidity (CHARM) program. *European journal of heart failure.* 2009;11(2):170-7. Epub 2009/01/27. doi: 10.1093/eurjhf/hfn031. PubMed PMID: 19168515; PubMed Central PMCID: PMCPMC2639422.
28. Moller S, Bernardi M. Interactions of the heart and the liver. *European heart journal.* 2013;34(36):2804-11. doi: 10.1093/eurheartj/ehs446. PubMed PMID: 23853073.

29. Nagy Á, Holndonner-Kirst E, Eke C, Szécsi B, Szabó A, Plamondon MJ, Perioperative Low Tetraiodothyronine Levels and Adverse Outcomes After Heart Transplantation: A Retrospective, Observational Study. *J Cardiothorac Vasc Anesth.* 2020;34(10):2648-54. Epub 20200421. doi: 10.1053/j.jvca.2020.03.052. PubMed PMID: 32389455.
30. Nagy A, Holndonner-Kirst E, Eke C, Kertai MD, Fazekas L, Benke K, Model for end-stage liver disease scores in veno-arterial extracorporeal membrane oxygenation. *Int J Artif Organs.* 2020;43(10):684-91. Epub 2020/02/27. doi: 10.1177/0391398820906538. PubMed PMID: 32098569.
31. Scheinfeld MH, Bilali A, Koenigsberg M. Understanding the spectral Doppler waveform of the hepatic veins in health and disease. *Radiographics.* 2009;29(7):2081-98. doi: 10.1148/rg.297095715. PubMed PMID: 19926763.
32. Abu-Yousef MM. Duplex Doppler sonography of the hepatic vein in tricuspid regurgitation. *AJR American journal of roentgenology.* 1991;156(1):79-83. doi: 10.2214/ajr.156.1.1898574. PubMed PMID: 1898574.
33. Yoshida K, Yoshikawa J, Kato H, Yanagihara K, Okumachi F, Koizumi K, [Tricuspid regurgitation evaluated by Doppler hepatic vein flow patterns]. *J Cardiogr.* 1985;15(4):1157-69. Epub 1985/12/01. PubMed PMID: 3879835.
34. Zabielska J, Zabielski G. [The usefulness of Duplex-Doppler sonography in estimation of hepatic veins and inferior vena cava in the group of patients treated for congestive heart disease]. *Pol Merkur Lekarski.* 2007;23(137):328-34. Epub 2008/03/26. PubMed PMID: 18361313.
35. Scheinfeld MH, Bilali A, Koenigsberg M. Understanding the Spectral Doppler Waveform of the Hepatic Veins in Health and Disease. *RadioGraphics.* 2009;29(7):2081-98. doi: 10.1148/rg.297095715. PubMed PMID: 19926763.
36. Fadel BM, Alassas K, Husain A, Dahdouh Z, Di Salvo G. Spectral Doppler of the Hepatic Veins in Noncardiac Diseases: What the Echocardiographer Should Know. *Echocardiography.* 2015;32(9):1424-7. doi: 10.1111/echo.12994.
37. Fadel BM, Almulla K, Husain A, Dahdouh Z, Di Salvo G, Mohty D. Spectral Doppler of the Hepatic Veins in Tricuspid Valve Disease. *Echocardiography.* 2015;32(5):856-9. doi: 10.1111/echo.12773.

38. Via G, Tavazzi G, Bendjelid K. The right ventricle after cardiopulmonary bypass: new insights on its adaptive physiology. *J Clin Monit Comput.* 2020;34(6):1133-4. Epub 2020/03/12. doi: 10.1007/s10877-020-00497-9. PubMed PMID: 32157492.
39. Korshin A, Gronlykke L, Holmgaard F, Kjoller SM, Gustafsson F, Nilsson JC, Right ventricular transverse displacement increases following cardiac surgery: possibly compensating loss in tricuspid annular plane systolic excursion (TAPSE). *J Clin Monit Comput.* 2020;34(6):1139-48. Epub 2020/01/27. doi: 10.1007/s10877-020-00466-2. PubMed PMID: 31983013.
40. Lim SY, Kim S. Pathophysiology of Cardiorenal Syndrome and Use of Diuretics and Ultrafiltration as Volume Control. *Korean Circ J.* 2021;51(8):656-67. Epub 2021/07/07. doi: 10.4070/kcj.2021.0996. PubMed PMID: 34227267; PubMed Central PMCID: PMCPCMC8326215.
41. Ueno K, Seki S, Shiokawa N, Matsuba T, Miyazono A, Hazeki D, Validation of acute kidney injury according to the modified KDIGO criteria in infants after cardiac surgery for congenital heart disease. *Nephrology (Carlton).* 2019;24(3):294-300. Epub 2018/02/17. doi: 10.1111/nep.13240. PubMed PMID: 29451341.
42. Machado MN, Nakazone MA, Maia LN. Acute kidney injury based on KDIGO (Kidney Disease Improving Global Outcomes) criteria in patients with elevated baseline serum creatinine undergoing cardiac surgery. *Rev Bras Cir Cardiovasc.* 2014;29(3):299-307. Epub 2014/11/06. doi: 10.5935/1678-9741.20140049. PubMed PMID: 25372901; PubMed Central PMCID: PMCPCMC4412317.
43. Harky A, Joshi M, Gupta S, Teoh WY, Gatta F, Snosi M. Acute Kidney Injury Associated with Cardiac Surgery: a Comprehensive Literature Review. *Braz J Cardiovasc Surg.* 2020;35(2):211-24. Epub 2020/05/06. doi: 10.21470/1678-9741-2019-0122. PubMed PMID: 32369303; PubMed Central PMCID: PMCPCMC7199993.
44. Rosner MH, Okusa MD. Acute kidney injury associated with cardiac surgery. *Clin J Am Soc Nephrol.* 2006;1(1):19-32. Epub 2007/08/21. doi: 10.2215/CJN.00240605. PubMed PMID: 17699187.
45. Mao H, Katz N, Ariyanon W, Blanca-Martos L, Adybelli Z, Giuliani A, Cardiac surgery-associated acute kidney injury. *Cardiorenal Med.* 2013;3(3):178-99. Epub 2014/01/24. doi: 10.1159/000353134. PubMed PMID: 24454314; PubMed Central PMCID: PMCPCMC3884176.

46. Cheungpasitporn W, Thongprayoon C, Kittanamongkolchai W, Srivali N, O'Corragain OA, Edmonds PJ, Comparison of renal outcomes in off-pump versus on-pump coronary artery bypass grafting: A systematic review and meta-analysis of randomized controlled trials. *Nephrology (Carlton)*. 2015;20(10):727-35. Epub 2015/05/15. doi: 10.1111/nep.12506. PubMed PMID: 25968971.
47. Lamy A, Devereaux PJ, Prabhakaran D, Taggart DP, Hu S, Paolasso E, Off-pump or on-pump coronary-artery bypass grafting at 30 days. *N Engl J Med*. 2012;366(16):1489-97. Epub 2012/03/28. doi: 10.1056/NEJMoa1200388. PubMed PMID: 22449296.
48. Bruins P, te Velthuis H, Yazdanbakhsh AP, Jansen PG, van Hardevelt FW, de Beaumont EM, Activation of the complement system during and after cardiopulmonary bypass surgery: postsurgery activation involves C-reactive protein and is associated with postoperative arrhythmia. *Circulation*. 1997;96(10):3542-8. Epub 1997/12/13. doi: 10.1161/01.cir.96.10.3542. PubMed PMID: 9396453.
49. Plestina S, Gamulin S. Kidney ischaemia-reperfusion injury and polyribosome structure. *Nephron*. 2001;89(2):201-7. Epub 2001/09/11. doi: 10.1159/000046068. PubMed PMID: 11549903.
50. Birnie K, Verheyden V, Pagano D, Bhabra M, Tilling K, Sterne JA, Predictive models for kidney disease: improving global outcomes (KDIGO) defined acute kidney injury in UK cardiac surgery. *Crit Care*. 2014;18(6):606. Epub 2015/02/13. doi: 10.1186/s13054-014-0606-x. PubMed PMID: 25673427; PubMed Central PMCID: PMC4258283.
51. Howitt SH, Grant SW, Caiado C, Carlson E, Kwon D, Dimarakis I, The KDIGO acute kidney injury guidelines for cardiac surgery patients in critical care: a validation study. *BMC Nephrol*. 2018;19(1):149. Epub 2018/06/27. doi: 10.1186/s12882-018-0946-x. PubMed PMID: 29940876; PubMed Central PMCID: PMC6020229.
52. Yu Y, Li C, Zhu S, Jin L, Hu Y, Ling X, Diagnosis, pathophysiology and preventive strategies for cardiac surgery-associated acute kidney injury: a narrative review. *Eur J Med Res*. 2023;28(1):45. Epub 2023/01/25. doi: 10.1186/s40001-023-00990-2. PubMed PMID: 36694233; PubMed Central PMCID: PMC9872411.
53. Messmer AS, Zingg C, Müller M, Gerber JL, Schefold JC, Pfortmueller CA. Fluid Overload and Mortality in Adult Critical Care Patients-A Systematic Review and Meta-

Analysis of Observational Studies. *Crit Care Med.* 2020;48(12):1862-70. doi: 10.1097/CCM.0000000000004617. PubMed PMID: 33009098.

54. Roy S, Couture P, Qizilbash B, Toupin F, Levesque S, Carrier M, Hemodynamic pressure waveform analysis in predicting fluid responsiveness. *J Cardiothorac Vasc Anesth.* 2013;27(4):676-80. doi: 10.1053/j.jvca.2012.11.002. PubMed PMID: 23849524.

55. Malbrain ML, Marik PE, Witters I, Cordemans C, Kirkpatrick AW, Roberts DJ, Fluid overload, de-resuscitation, and outcomes in critically ill or injured patients: a systematic review with suggestions for clinical practice. *Anaesthesiol Intensive Ther.* 2014;46(5):361-80. doi: 10.5603/AIT.2014.0060. PubMed PMID: 25432556.

56. Malbrain M, Martin G, Ostermann M. Everything you need to know about deresuscitation. *Intensive Care Med.* 2022. Epub 2022/08/07. doi: 10.1007/s00134-022-06761-7. PubMed PMID: 35932335; PubMed Central PMCID: PMCPMC9362613.

57. Pettey G, Hermansen JL, Nel S, Moutlana HJ, Muteba M, Juhl-Olsen P, Ultrasound Hepatic Vein Ratios Are Associated With the Development of Acute Kidney Injury After Cardiac Surgery. *J Cardiothorac Vasc Anesth.* 2021. Epub 2021/08/23. doi: 10.1053/j.jvca.2021.07.039. PubMed PMID: 34419361.

58. Barili F, Pacini D, D'Ovidio M, Dang NC, Alamanni F, Di Bartolomeo R, The Impact of EuroSCORE II Risk Factors on Prediction of Long-Term Mortality. *Ann Thorac Surg.* 2016;102(4):1296-303. Epub 2016/06/21. doi: 10.1016/j.athoracsur.2016.04.017. PubMed PMID: 27319983.

59. Nashef SA, Roques F, Michel P, Gauducheau E, Lemeshow S, Salamon R. European system for cardiac operative risk evaluation (EuroSCORE). *Eur J Cardiothorac Surg.* 1999;16(1):9-13. Epub 1999/08/24. doi: 10.1016/s1010-7940(99)00134-7. PubMed PMID: 10456395.

60. Abe S, Yoshihisa A, Takiguchi M, Shimizu T, Nakamura Y, Yamauchi H, Liver dysfunction assessed by model for end-stage liver disease excluding INR (MELD-XI) scoring system predicts adverse prognosis in heart failure. *PLoS One.* 2014;9(6):e100618. doi: 10.1371/journal.pone.0100618. PubMed PMID: 24955578; PubMed Central PMCID: PMC4067358.

61. Farr M, Mitchell J, Lippel M, Kato TS, Jin Z, Ippolito P, Combination of liver biopsy with MELD-XI scores for post-transplant outcome prediction in patients with advanced heart failure and suspected liver dysfunction. *The Journal of heart and lung*

transplantation : the official publication of the International Society for Heart Transplantation. 2015;34(7):873-82. doi: 10.1016/j.healun.2014.12.009. PubMed PMID: 25851466; PubMed Central PMCID: PMC4941637.

62. Thakar CV, Arrigain S, Worley S, Yared JP, Paganini EP. A clinical score to predict acute renal failure after cardiac surgery. *J Am Soc Nephrol.* 2005;16(1):162-8. Epub 20041124. doi: 10.1681/ASN.2004040331. PubMed PMID: 15563569.

63. Yokoyama H, Sekino M, Funaoka H, Sato S, Araki H, Egashira T, Association between enterocyte injury and fluid balance in patients with septic shock: a post hoc exploratory analysis of a prospective observational study. *BMC Anesthesiol.* 2021;21(1):293. Epub 2021/11/25. doi: 10.1186/s12871-021-01515-2. PubMed PMID: 34814831; PubMed Central PMCID: PMC8609797.

64. Yang W, Chan FP, Reddy VM, Marsden AL, Feinstein JA. Flow simulations and validation for the first cohort of patients undergoing the Y-graft Fontan procedure. *J Thorac Cardiovasc Surg.* 2015;149(1):247-55. Epub 2014/12/03. doi: 10.1016/j.jtcvs.2014.08.069. PubMed PMID: 25439766.

65. Rosenkranz S, Howard LS, Gombert-Maitland M, Hoepfer MM. Systemic Consequences of Pulmonary Hypertension and Right-Sided Heart Failure. *Circulation.* 2020;141(8):678-93. Epub 20200224. doi: 10.1161/CIRCULATIONAHA.116.022362. PubMed PMID: 32091921.

66. Verbrugge FH, Grieten L, Mullens W. Management of the cardiorenal syndrome in decompensated heart failure. *Cardiorenal Med.* 2014;4(3-4):176-88. doi: 10.1159/000366168. PubMed PMID: 25737682; PubMed Central PMCID: PMC4299260.

67. Iida N, Seo Y, Sai S, Machino-Ohtsuka T, Yamamoto M, Ishizu T, Clinical Implications of Intrarenal Hemodynamic Evaluation by Doppler Ultrasonography in Heart Failure. *JACC Heart Fail.* 2016;4(8):674-82. Epub 2016/05/18. doi: 10.1016/j.jchf.2016.03.016. PubMed PMID: 27179835.

68. Monnet X, Dres M, Ferre A, Le Teuff G, Jozwiak M, Bleibtreu A, Prediction of fluid responsiveness by a continuous non-invasive assessment of arterial pressure in critically ill patients: comparison with four other dynamic indices. *Br J Anaesth.* 2012;109(3):330-8. Epub 2012/06/28. doi: 10.1093/bja/aes182. PubMed PMID: 22735299.

69. Monnet X, Malbrain M, Pinsky MR. The prediction of fluid responsiveness. *Intensive Care Med.* 2023;49(1):83-6. Epub 2022/11/04. doi: 10.1007/s00134-022-06900-0. PubMed PMID: 36323911.
70. Monnet X, Marik PE, Teboul JL. Prediction of fluid responsiveness: an update. *Ann Intensive Care.* 2016;6(1):111. Epub 2016/11/20. doi: 10.1186/s13613-016-0216-7. PubMed PMID: 27858374; PubMed Central PMCID: PMC5114218.
71. Jozwiak M, Monnet X, Teboul JL. Prediction of fluid responsiveness in ventilated patients. *Ann Transl Med.* 2018;6(18):352. Epub 2018/10/30. doi: 10.21037/atm.2018.05.03. PubMed PMID: 30370279; PubMed Central PMCID: PMC6186547.
72. Huang CC, Fu JY, Hu HC, Kao KC, Chen NH, Hsieh MJ, Prediction of fluid responsiveness in acute respiratory distress syndrome patients ventilated with low tidal volume and high positive end-expiratory pressure. *Crit Care Med.* 2008;36(10):2810-6. Epub 2008/09/04. doi: 10.1097/CCM.0b013e318186b74e. PubMed PMID: 18766099.
73. Huang D, Ma J, Wang S, Qin T, Song F, Hou T, Changes of cardiac output and velocity time integral in blood return at the end of renal replacement therapy predict fluid responsiveness in critically ill patients with acute circulatory failure. *BMC Anesthesiol.* 2023;23(1):25. Epub 2023/01/14. doi: 10.1186/s12871-023-01976-7. PubMed PMID: 36639628; PubMed Central PMCID: PMC9840273.
74. Guerin L, Monnet X, Teboul JL. Monitoring volume and fluid responsiveness: from static to dynamic indicators. *Best Pract Res Clin Anaesthesiol.* 2013;27(2):177-85. Epub 2013/09/10. doi: 10.1016/j.bpa.2013.06.002. PubMed PMID: 24012230.
75. Gan H, Cannesson M, Chandler JR, Ansermino JM. Predicting fluid responsiveness in children: a systematic review. *Anesth Analg.* 2013;117(6):1380-92. Epub 2013/11/22. doi: 10.1213/ANE.0b013e3182a9557e. PubMed PMID: 24257389.
76. Sugawara Y, Yoshihisa A, Ishibashi S, Matsuda M, Yamadera Y, Ohara H, Liver Congestion Assessed by Hepatic Vein Waveforms in Patients With Heart Failure. *CJC Open.* 2021;3(6):778-86. Epub 2021/06/26. doi: 10.1016/j.cjco.2021.02.004. PubMed PMID: 34169257; PubMed Central PMCID: PMC8209365.
77. Guerrero Gomez A, Gonzalez Jaramillo N, Castro Perez JA. Ultra-fast-track extubation vs. conventional extubation after cardiac surgery in a cardiovascular reference centre in Colombia. A longitudinal study. *Rev Esp Anesthesiol Reanim (Engl Ed).*

2019;66(1):10-7. Epub 2018/07/29. doi: 10.1016/j.redar.2018.06.005. PubMed PMID: 30054093.

78. Ellenbroek DFJ, van Kessel L, Compagner W, Brouwer T, Bouwman RA, van Straten B, Diagnostic performance of echocardiography to predict cardiac tamponade after cardiac surgery. *Eur J Cardiothorac Surg.* 2022;62(1). Epub 2021/11/19. doi: 10.1093/ejcts/ezab468. PubMed PMID: 34791128; PubMed Central PMCID: PMC9257667.

79. Mandoli GE, Cameli M, Novo G, Agricola E, Righini FM, Santoro C, Right ventricular function after cardiac surgery: the diagnostic and prognostic role of echocardiography. *Heart Fail Rev.* 2019;24(5):625-35. Epub 2019/04/15. doi: 10.1007/s10741-019-09785-2. PubMed PMID: 30982175.

80. Heiberg J, El-Ansary D, Royse CF, Royse AG, Alsaddique AA, Canty DJ. Transthoracic and transoesophageal echocardiography: a systematic review of feasibility and impact on diagnosis, management and outcome after cardiac surgery. *Anaesthesia.* 2016;71(10):1210-21. Epub 2016/06/28. doi: 10.1111/anae.13545. PubMed PMID: 27341788.

81. Carricart M, Denault AY, Couture P, Limoges P, Babin D, Levesque S, Incidence and significance of abnormal hepatic venous Doppler flow velocities before cardiac surgery. *J Cardiothorac Vasc Anesth.* 2005;19(6):751-8. Epub 2005/12/06. doi: 10.1053/j.jvca.2004.11.052. PubMed PMID: 16326300.

82. Caraba A, Iurciuc S, Munteanu A, Iurciuc M. Hyponatremia and Renal Venous Congestion in Heart Failure Patients. *Dis Markers.* 2021;2021:6499346. Epub 2021/08/24. doi: 10.1155/2021/6499346. PubMed PMID: 34422137; PubMed Central PMCID: PMC9257667.

83. Sandrikov VA, Dzeranova AN, Fedulova SV, Lokshin LS, Karshieva AR, Kulagina TY. [ASSESSMENT OF LIVER FUNCTION WITH TRANSESOPHAGEAL ECHOCARDIOGRAPHY HEART SURGERY WITH CARDIOPULMONARY BYPASS]. *Anesteziol Reanimatol.* 2016;61(1):4-7. Epub 2016/05/20. PubMed PMID: 27192845.

84. Petaja L, Vaara S, Liuhanen S, Suojaranta-Ylinen R, Mildh L, Nisula S, Acute Kidney Injury After Cardiac Surgery by Complete KDIGO Criteria Predicts Increased

- Mortality. *J Cardiothorac Vasc Anesth.* 2017;31(3):827-36. Epub 2016/11/20. doi: 10.1053/j.jvca.2016.08.026. PubMed PMID: 27856153.
85. Giallourakis CC, Rosenberg PM, Friedman LS. The liver in heart failure. *Clin Liver Dis.* 2002;6(4):947-67, viii-ix. Epub 2003/01/09. doi: 10.1016/s1089-3261(02)00056-9. PubMed PMID: 12516201.
86. Okuda H, Ajima T, Obata H. [Severe heart failure and liver dysfunction]. *Kokyu to junkan Respiration & circulation.* 1992;40(3):233-8. PubMed PMID: 1579743.
87. Fuhrmann V, Kneidinger N, Herkner H, Heinz G, Nikfardjam M, Bojic A, Hypoxic hepatitis: underlying conditions and risk factors for mortality in critically ill patients. *Intensive Care Med.* 2009;35(8):1397-405. doi: 10.1007/s00134-009-1508-2. PubMed PMID: 19506833.
88. Nikolaou M, Parissis J, Yilmaz MB, Seronde MF, Kivikko M, Laribi S, Liver function abnormalities, clinical profile, and outcome in acute decompensated heart failure. *European heart journal.* 2013;34(10):742-9. doi: 10.1093/eurheartj/ehs332. PubMed PMID: 23091203.
89. Parissis J, Nikolaou M, Mebazaa A. The Model for End-stage Liver Disease score in acute heart failure: hepatorenal dysfunction hides behind. *European journal of heart failure.* 2016;18(12):1522-3. doi: 10.1002/ejhf.681. PubMed PMID: 27813334.

## 12. Bibliography of the candidate's publications

### **Related to thesis:**

Eke, Csaba ; Szabó, András ; Nagy, Ádám ; Szécsi, Balázs ; Szentgróti, Rita ; Dénes, András ; Kertai, Miklós D. ; Fazekas, Levente ; Kovács, Attila ; Lakatos, Bálint Merkely, Béla ; Székely, Andrea

Association between Hepatic Venous Congestion and Adverse Outcomes after Cardiac Surgery

DIAGNOSTICS 12 : 12 Paper: 3175 , 12 p. (2022)

IF: 3,6

Eke, Csaba ; Szabó, András ; Nagy, Ádám ; Párkányi, Boglár ; Kertai, Miklós D. ; Fazekas, Levente ; Kovács, Attila ; Lakatos, Bálint ; Hartyánszky, István ; Gál, János Merkely, Béla ; Székely, Andrea

Association between Preoperative Retrograde Hepatic Vein Flow and Acute Kidney Injury after Cardiac Surgery

DIAGNOSTICS 12 : 3 Paper: 699 , 11 p. (2022)

IF: 3,6

### **Not related to thesis:**

Szabó, András ✉ ; Szabó, Dominika ; Tóth, Krisztina ; Szécsi, Balázs ; Szentgróti, Rita ; Nagy, Ádám ; Eke, Csaba ; Sándor, Ágnes ; Benke, Kálmán ; Merkely, Béla

Comprehensive frailty assessment with multidimensional frailty domains as a predictor of mortality among vascular and cardiac surgical patients

PHYSIOLOGY INTERNATIONAL 110 : 2 pp. 191-210. , 20 p. (2023)

IF: 1,4

Tóth, Krisztina ; Szabó, András ; Nagy, Ádám ; Szabó, Dominika ; Szécsi, Balázs ; Eke, Csaba ; Sándor, Ágnes ; Susánszky, Éva ; Holndonner-Kirst, Enikő ; Merkely, Béla  
Preoperative nutritional state is associated with mid- and long-term mortality after cardiac surgery  
ANNALS OF PALLIATIVE MEDICINE 10 : 11 pp. 11333-11347. , 15 p. (2021)  
IF: 0,53

Szabo, Andras ; Szabo, Dominika ; Toth, Krisztina ; Szecsi, Balazs ; Sandor, Anges ; Eke, Csaba ; Szentgroti, Rita ; Parkanyi, Boglar ; Denes, Andras ; Szekely, Andrea  
EARLY POSTOPERATIVE SERUM OSMOLARITY IMBALANCE MAY PREDICT WORSE SHORT-TERM MORTALITY AFTER HEART TRANSPLANTATION IN ADULTS  
JOURNAL OF CARDIOTHORACIC AND VASCULAR ANESTHESIA 35 : Suppl. 1 pp. S41-S41. Paper: PP:27 (2021)  
IF: 2,8

Nagy, Á. ; Szécsi, B.\* ; Eke, C. ; Szabó, A. ; Mihály, S. ; Fazekas, L. ; Hartyánszky, I. ; Párkányi, B. ; Holndonner-Kirst, E. ; Lex, D.  
Endocrine Management and Hormone Replacement Therapy in Cardiac Donor Management: A Retrospective Observational Study  
TRANSPLANTATION PROCEEDINGS 53 : 10 pp. 2807-2815. , 9 p. (2021)  
IF: 0,29

Moskola, Vladimír ✉ ; Eke, Csaba ; Takács, Johanna ; Susánszky, Éva ; Székely, Andrea ; Hornyák, István ; Ozsvárt, Barbara ; Néninger, Tibor ; Balogh, Zoltán  
Investigating the attitude of ambulance workers towards death  
KONTAKT 23 : 1 pp. 14-19. , 6 p. (2021)  
IF: 0,3

Szabó, András ✉; Tóth, Krisztina; Nagy, Ádám; Domokos, Dominika; Czobor, Nikoletta; Eke, Csaba; Sándor, Ágnes; Merkely, Béla; Susánszky, Éva; Gál, János  
The effect of cognitive dysfunction on mid- and long-term mortality after vascular surgery  
BMC GERIATRICS 21 : 1 Paper: 46 , 11 p. (2021)  
IF: 4,1

Nagy, Ádám; Holndonner-Kirst, Enikő; Eke, Csaba; Szécsi, Balázs; Szabó, András; Plamondon, Marie-Jo; Fazekas, Levente; Polos, Miklós; Benke, Kálmán; Szabolcs, Zoltán  
Perioperative Low Tetraiodothyronine Levels and Adverse Outcomes After Heart Transplantation: A Retrospective, Observational Study  
JOURNAL OF CARDIOTHORACIC AND VASCULAR ANESTHESIA 34 : 10 pp. 2648-2654. , 7 p. (2020)  
IF: 2,8

Nagy, Ádám; Holndonner-Kirst, Enikő; Eke, Csaba; Kertai, Miklós D; Fazekas, Levente; Benke, Kálmán; Pólos, Miklós; Szabolcs, Zoltán; Hartyánszky, István; Gál, János  
Model for end-stage liver disease scores in veno-arterial extracorporeal membrane oxygenation  
INTERNATIONAL JOURNAL OF ARTIFICIAL ORGANS 43 : 10 pp. 684-691. , 8 p. (2020)  
IF: 1,7

Nagy, A; Szécsi, B; Lendvai, M; Eke, Cs; Holndonner-Kirst, E; Szabó, A; Czobor, N; Gál, J; Merkely, B; Székely, A  
Perioperative thyroid dysfunction and outcome after heart transplantation  
JOURNAL OF CARDIOTHORACIC AND VASCULAR ANESTHESIA 33 : Suppl. 2 pp. S159-S159. , 1 p. (2019)  
IF: 2,8

### 13. Acknowledgements

I would like to thank my supervisor, Prof. Andrea Székely for all her help in the progress of my research and clinical career and the lots of good advice. Thank you for giving me a chance while I was a medical student.

I am grateful for Ádám Nagy MD, to help me get started in my science work when I was still a medical student.

I would like to thank my friend Krisztina Tóth MD, for the common work and experiences.

I would like to thank my family and friends all the emotional support they gave me during my studies, András Szilvay, Zsolt Benkő and Dávid Karádi for always being there for me, my mother and grandmother for the stable background.

I appreciate the opportunity to conduct our studies in the Department of Cardiac Surgery of the Heart and Vascular Centre, Semmelweis University.

I am grateful to all the cardiac surgery patients who agreed to participate in our research between 2018 and 2022, helping us to advance our scientific knowledge.

Finally, everyone deserves to be thanked, who I don't have a chance to mention by name, but was involved on the road to my Ph.D.